

Novel solutions of RG equations for $\alpha(s)$ and $\beta(\alpha)$ in the large N_c limit

Yu.A. Simonov

Abstract

General solution of RG equations in the framework of background perturbation theory is written down in the large N_c limit. A simplified (model) approximation to the general solution is suggested which allows to write $\beta(\alpha)$ and $\alpha(\beta)$ to any loop order. The resulting $\alpha_B(Q^2)$ coincides asymptotically at large $|Q^2|$ with standard (free) α_s , saturates at small $Q^2 \geq 0$ and has poles at time-like Q^2 in agreement with analytic properties of physical amplitudes in the large N_c limit.

1. The running α_s of Standard Perturbation Theory (SPT) has unphysical singularities (Landau ghost pole and cuts) in the Euclidean region; which contradict analyticity of physical amplitudes and are not observed in lattice calculations of $\alpha_s(Q^2)$ [1].

As it was shown in [2, 3] the Background Perturbation Theory (BPT) is free of these defects and the coupling constant α_B of BPT displays an important property of IR freezing (saturation) with $\alpha_B(Q^2 = 0) \approx 0.5$ [2, 3]. This behaviour of α_B is well confirmed by experimental data on spin splitting of quarkonia levels [4, 5] and by lattice data [6].

Analytic properties of $\alpha_B(Q^2)$ in the Q^2 plane were not studied heretofore, and is the topic of the present letter. To this end the large N_c limit is exploited which ensures that any physical amplitude has only simple poles [7]. This requires that $\alpha_B(Q^2)$ should be a meromorphic function of Q^2 .

To find it explicitly we study below the general solution of RG equations and suggest a simple model approximation to it, yielding $\beta(\alpha)$ to all orders and explicit form of $\alpha_B(Q^2)$.

Resulting $\alpha_B(Q^2)$ can be compared with $\alpha_s(Q^2)$ of SPT and coincides with the latter asymptotically at large Q^2 .

2. Separating background field B_μ in the total gluonic field $A_\mu = B_\mu + a_\mu$ and expanding in ga_μ one gets the BPT series, where coefficients depend both on external momenta and the background field gB_μ , which turns out to be RG invariant, if background gauge is chosen [8]. Correspondingly $\alpha_B \equiv g^2/4\pi$ satisfies the standard RG equations, where momenta and averaged background characteristic $\text{tr} < F_{\mu_1\nu_1}(B)F_{\mu_2\nu_2}(B)\dots F_{\mu_n\nu_n}(B) >$ enter together and will be denoted $\{\mathcal{P}_i^2\}$.

One starts with the standard definition

$$\frac{d\alpha_B(\mu)}{d\ln\mu} = \beta(\alpha_B) \quad (1)$$

and represents $\beta(\alpha_B)$ in the form

$$\beta(\alpha) = -\frac{\beta_0}{2\pi} \frac{\alpha^2}{[1 - \frac{\beta_1}{2\pi\beta_0}\varphi'(\frac{1}{\alpha})]} \quad (2)$$

where $\varphi(z)$ is an arbitrary function with some conditions to be imposed below, and prime means derivative.

Solving (1) with the help of (2) one has

$$\alpha_B = \frac{4\pi}{\beta[\ln\mu^2 + \chi + \frac{2\beta_1}{\beta_0^2}\varphi(\frac{1}{\alpha_B})]} \quad (3)$$

Here χ is an arbitrary dimensionless function of $\{\mathcal{P}_i^2\}$, while φ is partly fixed by few coefficients of expansion of $\beta(\alpha)$,

$$\beta(\alpha) = -\frac{\beta_0}{2\pi}\alpha^2 - \frac{\beta_1}{4\pi^2}\alpha^3 - \frac{\beta_2}{64\pi^3}\alpha^4 - \frac{\beta_3\alpha^5}{(4\pi)^4} \quad (4)$$

(Note that β_i are defined as in [9], for $n_f = 0, N_c = 3$ and $\overline{MS}$ scheme $\beta_0 = 11, \beta_1 = 51, \beta_2 = 2857, \beta_3 = 58486$.)

At this point we shall use the large N_c limit and require that $\alpha_B(Q^2)$ be a meromorphic function of Q^2 . Consider e.g. the process e^+e^- into hadrons, and the photon self-energy part $\Pi(Q^2)$, which has a pole expansion [3]

$$\Pi(Q^2, \alpha_B) = \sum_{n=0}^{\infty} \frac{C_n(\alpha_B)}{Q^2 + M_n^2(\alpha_B)} + \text{subtractions} \quad (5)$$

Expanding in α_B and doing a partial summation (equivalent to the summation of leading logarithms) one obtains a running $\alpha_B(Q^2)$ which has poles at some generally speaking, shifted positions as compared to the poles of the lowest order $-M_n^2(0)$. The latter are proportional to n ; $M_n^2(0) = m_0^2 n + M_0^2$; $n = 0, 1, 2, \dots$, $m_0^2 = 4\pi\sigma$, $C_n(0) = m_0^2$ [3]. The lowest (partonic) approximation thus reduces to $\Pi(Q^2, 0) \sim \psi(\frac{Q^2 + M_0^2}{m^2})$, where $\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$.

To specify the form of the function $\varphi(\frac{1}{\alpha_B})$ and $\chi(Q^2)$, we shall use two requirements:

- i) the correspondence principle with the SPT, which tells that for $Q^2 \gg \kappa^2$, where κ^2 is the scale of nonperturbative vacuum fields, our solution should coincide with that of SPT.
- ii) $\alpha_B(Q^2)$ should be meromorphic function of Q^2 , and hence both χ and φ should be meromorphic functions of their arguments (Q^2 and $\frac{1}{\alpha_B}$ respectively).

The first requirement means that coefficients C_n in the sum over poles should have finite limit C_∞ , i.e. one can represent this sum as the Euler function $\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$ plus a sum with fast decreasing coefficients, which we write symbolically as a finite sum. Thus the general solution satisfying i) and ii) can be written as

$$\ln \mu^2 + \chi = \ln \frac{m^2}{\Lambda^2} + \psi\left(\frac{Q^2 + M_0^2}{m^2}\right) + \sum_{n=1}^{N_1} \frac{b_n}{Q^2 + m_n^2} \quad (6)$$

$$\varphi\left(\frac{1}{\alpha_B}\right) = \psi\left(\frac{\lambda}{\alpha_B} + \Delta\right) + P\left(\frac{1}{\alpha_B}\right) \quad (7)$$

where Δ is some constant and $\lambda = 4\pi/\beta_0$. Note that our solution (6), (7) satisfies (1) and reproduces the first two coefficients in the expansion (4) when all $b_n = 0$ and $P \equiv 0$, while our β_2, β_3 appear to be smaller; however the latter are scheme-dependent. It is easy to reproduce β_2, β_3 by choosing in (7) $P \neq 0$. This choice of $\varphi(\frac{1}{\alpha_B})$ leads to a small numerical (around 2%) change in α_B as compared to the minimal choice (8) for all $Q^2 \leq 0$.

3. In what follows we confine ourselves to the minimal model of (6), (7) with all b_n , and P identically zero, which yields

$$\alpha_B = \frac{4\pi}{\beta_0} \left[\ln \frac{m^2}{\Lambda^2} + \psi\left(\frac{Q^2 + M_0^2}{m^2}\right) + \frac{2\beta_1}{\beta_0^2} \psi\left(\frac{\lambda}{\alpha_B} + \Delta\right) \right]^{-1} \quad (8)$$

Eq.(8) defines α_B as a meromorphic function of Q^2 in the whole Q^2 complex plane.

Moreover, Eq.(8) defines α_B to all orders in the “loop expansion” obtained by iteration of the last term in the denominator of (8) and will be compared below to the SPT loop expansion. Consider e.g. large Q^2 , $\frac{Q^2 + M_0^2}{m^2} \gg 1$. Then using the asymptotic expansion for $z \rightarrow \infty$

$$\psi(z) = \ln z - \frac{1}{2z} - \sum_{k=1}^{\infty} \frac{B_{2k}}{2kz^{2k}} \quad (9)$$

where B_n are Bernoulli numbers, one obtains to the lowest order

$$\alpha_B^{(0)} = \frac{4\pi}{\beta_0 \ln\left(\frac{Q^2 + M_0^2}{\Lambda^2}\right)} \quad (10)$$

which explicitly shows the AF behaviour for large Q^2 and absence of Landau pole (for $M_0^2 > \Lambda^2$).

The forms (2) and (8) admit in general poles of $\alpha_B(Q^2)$ for $Q^2 \geq 0$ and poles of $\beta(\alpha_B)$ for $\alpha_B \geq 0$, which would be unphysical. The latter poles are defined by equation $1 = \frac{\beta_1 \lambda}{2\pi \beta_0} \psi'\left(\frac{\lambda}{\alpha} + \Delta\right)$,

and are excluded from the region $\alpha \geq 0$ when $\Delta > \Delta_0 = 1.255$. This latter condition also excludes singularities of $\alpha_B(Q^2)$ (8), at $Q^2 > 0$. At the same time $\beta(\alpha)$ has a point of condensing zeros on the negative side of $\alpha = 0$, and $\beta(\alpha)$ is monotonically decreasing when $\alpha \rightarrow \infty$, as it is shown in Table 1. Note that a pole of $\beta(\alpha)$ at $\alpha = \alpha^* > 0$ leads to a situation when real solutions $\alpha(q)$ exist only for $\alpha \leq \alpha^*, Q > Q^*$.




Figure 1: Exact solution of Eq.(8) with $\Delta = 1.5$ as a function of $s \equiv Q^2$ (solid line), and exact solution for α_B from the extended model, Eq.(3) with φ from (21) and $\eta = 3, \Delta = 1.5$ which reproduces all known β_n coefficients (broken line).

4. Although $\alpha_B(Q^2)$ does not have poles for $\Delta > \Delta_0$ in the Euclidean region, $Q^2 > 0$, it has an infinite number of poles for $Q^2 < 0$.

These poles are given by zeros of the denominator in (8), which can be written as

$$-Q^2 \equiv s_n = s_n^{(0)} - \delta_n, \quad s_n^{(0)} \equiv M_n^2 = M_0^2 + nm^2 \quad (11)$$

and δ is then found from (8) to be [10]

$$\delta_n^{-1} \cong \ln \frac{M_n^2}{\Lambda^2}. \quad (12)$$

Hence poles of α_B are shifted as compared with poles of $\psi\left(\frac{Q^2 + M_0^2}{m^2}\right)$, the latter describe the physical states of (double) hybrids - the background analog of gluon loops renormalizing α_s in SPT. In the vicinity of the pole α_B has the form [10]

$$\alpha_B(s \approx s_n - \delta_n) = \frac{(4\pi/\beta_0 - \psi'(\Delta))}{\psi'(z_n)(z - z_n)}, \quad z = \frac{M_0^2 - s}{m^2} \quad (13)$$

Hence perturbative series of BPT is a sum over poles of α_B and nonperturbative poles of $\Pi(Q^2, 0)$ in (5) (dual to partonic contribution), all poles being in the region $s > M_0^2 - \delta_0 > 0$ and this situation is in accord with the large N_c analytic properties. Now one has to show that the

”physically averaged value” of $\alpha_B(s)$ corresponds to the AF behaviour, as it is seen e.g. in the e^+e^- annihilation. To this end one can choose two equivalent procedures: i) to introduce the width of the pole in $\alpha_B(s)$, or ii) equivalently to consider the BPT series at the shifted value of s , $s = \bar{s}(1 + i\gamma)$ (see e.g. [11]).

In what follows we shall exploit the second path. Considering $\text{Im}\alpha_B$ above the real axis, at $\bar{s}(1 + i\gamma)$ with $\bar{s}$ large and γ fixed $\gamma \ll \pi$, allows to use asymptotics (10) (in one loop order) with the result

$$\text{Im}\alpha(\bar{s}(1 + i\gamma)) \approx \frac{4\pi(\pi - \gamma)}{b_0[\ln^2|\frac{M_0^2 - \bar{s}}{\Lambda^2}| + (\pi - \gamma)^2]} \quad (14)$$

Another form of discontinuity results from the expansion of the Adler function $D(Q^2) = Q^2 \frac{d\Pi(Q^2)}{d\ln Q^2}$, from which the hadronic ratio $R(s)$ can be obtained as [12].

$$R(s) = \frac{1}{2\pi i} \int_{-s-i\varepsilon}^{-s+i\varepsilon} D(\sigma) \frac{d\sigma}{\sigma} = \frac{1}{2\pi i} \int_{C(s)} D(\sigma) \frac{d\sigma}{\sigma} \quad (15)$$

Here $C(s)$ is the circle of radius s around the origin, comprising all singularities of $D(\sigma)$ in Euclidean region.

Similarly to [12] one can define the operation $\Phi\{\frac{\alpha_B}{\pi}\}$ transforming the known perturbative series for $R(s)$;

$$R(s) = N_c \sum_q e_q^2 \left[1 + \bar{\Phi}\left\{\frac{\alpha_B}{\pi}\right\} + d_2 \bar{\Phi}\left\{\left(\frac{\alpha_B}{\pi}\right)^2\right\} \right] \quad (16)$$

where

$$\bar{\Phi}\left\{\left(\frac{\alpha_B}{\pi}\right)^k\right\} = \frac{1}{2\pi i} \int_{C(s)} \left(\frac{\alpha_B(\sigma)}{\pi}\right)^k \frac{d\sigma}{\sigma} \quad (17)$$

We shall consider the situation, when one can use the logarithmic asymptotics (10) for α_B in the Minkowskian region. This happens in the physical situation described above: when poles have imaginary parts (widths of resonances), or equivalently when one consider effective $\bar{R}(s) = \frac{1}{2i}(\Pi(s(1 + i\gamma)) - \Pi(s(1 - i\gamma)))$ with fixed γ (so that requirement $|\arg(z = \frac{M_0^2 - s}{m^2})| < \pi$ is fulfilled).

In this case one obtains, inserting in the integral (17) the asymptotic form (10) (see [10] for details)

$$\Phi\left\{\frac{\bar{\alpha}_B}{\pi}\right\} \cong \frac{4}{b_0} \left\{ \frac{1}{\pi} \text{arctg} \frac{\pi}{\ln s/\Lambda^2} + \frac{2M_0^2}{s \ln^3 s/\Lambda^2} + O\left(\frac{M_0^2}{s \ln^4 s/\Lambda^2}\right) \right\} \quad (18)$$

5. One can now compare the power (“loop expansion”) of α_B in (8) in powers of $\alpha_B^{(0)}$ (10) with the SPT expression for α_s [9, 13]. To this end we shall write it as follows:

$$\alpha_B = \alpha_B^{(0)} \left\{ 1 - \frac{2\beta_1 \ln L}{\beta_0^2 L} + \frac{4\beta_1^2}{\beta_0^4 L^2} \left[\left(\ln L - \frac{1}{2} \right)^2 + b \right] \right\} \quad (19)$$

where $L \equiv \ln(\frac{Q^2 + M_0^2}{\Lambda^2})$ and for $n_f = 0$, $\Delta = 1.5$

$$b = -\frac{\beta_0^2}{2\beta_1} \left(\Delta - \frac{1}{2} \right) - \frac{1}{4} = -1.436 \quad (20)$$

This should be compared with the $\overline{MS}$ value of SPT, $b_{\overline{MS}} = 0.26$ ($n_f = 0$).

To bring our theoretical expressions for β_2, β_3 in agreement with the computed $\overline{MS}$ values (see [9] for the corresponding references), one should generalize the minimal model considered above by adding a term to $\varphi(\frac{1}{\alpha})$:

$$\varphi'(\frac{1}{\alpha}) = \lambda \psi'(\frac{\lambda}{\alpha} + \Delta) + \frac{\tilde{C}\alpha^2}{1 + \alpha^2 \eta^2} \quad (21)$$

Choosing $\tilde{C} = 1.253$ and arbitrary real η , one obtains both β_2, β_3 in close agreement (within 1%) with $\overline{MS}$ values. In this case also b in (20) gets contribution from the new term and becomes $\tilde{b} = b + \frac{2\pi\beta_0}{\beta_1}\tilde{C} = 0.263$, which agrees with the SPT [9]. We study numerically both minimal model and the corrected one as explained above and present the results for $\beta(\alpha)$ in Table 1 and for $\alpha_B(Q^2)$ in Table 2.

From Table 1 it is clear that nonperturbative $\beta(\alpha)$, namely $\beta_{min}(\alpha)$, calculated as in (2), (7) with $P \equiv 0$ and $\beta_{ext}(\alpha)$ calculated as in (2), (21), both decrease more moderately (as $-\text{const } \alpha^2$) at $\alpha > 1$ as compared to purely perturbative 3 loop expression (4). This might indicate that the series (4) is asymptotic and should be cut off after few first terms (note that β_3 is larger than previous three coefficients).

For $Q^2 = (10 \text{ GeV})^2$ and $M_0 = 1 \text{ GeV}$ the correction in $\alpha_B(Q^2)$ introduced by replacements (21) is around 2% and it decreases for Q^2 growing.

Defining $\Lambda \equiv \Lambda_B = \Lambda_{\overline{MS}}$ (since α_B and α_s practically coincide for $Q^2 > 40 (\text{GeV})^2$) one can compare exact $\alpha_B(Q^2)$ obtained by solving (8) with $\alpha_s(Q^2)$ (two loop) of SPT and observe a significant difference only at small Q^2 , $Q^2 \leq 10 \text{ GeV}^2$.

One can see in Table 2 that the difference of $\alpha_B^{(min)}$ and $\alpha_s^{(2loop)}$ is 20% at 1 GeV^2 and drops to 10% at 5 GeV^2 , which implies that the use of $\alpha_B^{(min)}$ instead of α_s in the Euclidean region would not produce much change in the region $Q^2 > 1 \text{ GeV}^2$ while it improves the situation for $Q^2 \leq 1 \text{ GeV}^2$, where $\alpha_B^{(min)}$ saturates in agreement with experiment and lattice data (see [1]-[6] and [14] for the discussion of this point).




Figure 2: $\beta_{min}(\alpha)$ in the minimal model Eqs.(2), (7) with $P \equiv 0, \Delta = 1.5$, reproducing scheme-independent coefficients β_0, β_1 —dashed line, and $\beta_{ext}(\alpha)$ in the extended model Eqs.(2), (21) with $\varepsilon = 3$ and $\Delta = 1.5$ (solid line) and $\Delta = 3$ (dotted line).

6. We have used the large N_c limit to simplify analysis of analytic properties of background coupling constant α_B . This has enabled us to study $\alpha_B(Q^2)$ also for time-like Q^2 and to suggest a simplified model solution of exact RG equations, which yields α_B to all orders and prescribes the nonperturbative expression for $\beta(\alpha)$.

The resulting solution for α_B coincides with the standard α_s at large $|Q^2|$ and displays familiar AF properties. At the same time $\alpha_B(Q^2)$ stays finite everywhere in the Euclidean region,

demonstrating phenomenon of IR saturation. For time-like Q^2 the model $\alpha_B(Q^2)$ in the large N_c limit acquires pole singularities, which have been treated above in the averaging procedure. It is worth noting that our minimal model correctly reproduces both scheme-independent coefficients β_0, β_1 . At the same time scheme-dependent coefficients β_2, β_3 etc. are defined by additional pole terms with arbitrary constants in them. Several improvements are possible with the suggested model solution. First, one can consider $n_f \neq 0$, N_c finite, in which case poles would acquire width. This will not change much the averaged values of $\alpha_B(s)$ calculated above, but will enable one to make local predictions of resonance behaviour. Secondly, our model is too simplified, since $\psi(\frac{Q^2+M_0^2}{m^2})$ appearing in (8), is responsible only for the equidistant spectrum of hybrids, (equivalent of gluon loops), but this spectrum should not be equidistant for lowest states, which yields power corrections, considered in [10].

The author is grateful to N.O.Agasian and A.V.Radyushkin for helpful discussions; a partial financial support of RFFI grants, 00-02-17836, 00-15-96786 and INTAS grants 00-00110 and 00-00366 are gratefully acknowledged.

α	0	0.1	0.2	0.5	1	1.2	1.5	2
$\beta_{min}(\alpha)$	0	-0.0187	-0.08	-0.587	-2.85	-4.36	-7.38	-14.59
$\beta_{ext}(\alpha)$	0	-0.019	-0.082	-0.65	-3.36	-5.23	-9.04	-18.43
$\beta_{\overline{MS}}^{(3loop)}$	0	-0.01895	-0.0833	-0.767	-6.82	-13.56	-33.35	-115.24

Table 1: The QCD β -function $\beta(\alpha)$ for $n_f = 0$, $N_c = 3$ obtained numerically to all orders in α for the minimal model $\beta_{min}(\alpha)$ Eq.(2) with $\varphi = \frac{\lambda}{\alpha} + \Delta$, and $\Delta = 1.5$; $\lambda = 1.14$, and for the extended model β_{ext} with φ' from (21). The bottom line refers to $\beta(\alpha)$ in $\overline{MS}$ as given in Eq. (4).

Q^2 (GeV ²)	0	0.5	1	1.5	2	3	5	10	40	100
$\alpha_B^{(min)}$	0.546	0.394	0.337	0.306	0.286	0.26	0.232	0.201	0.159	0.139
$\alpha_B^{(ext)}$	0.754	0.462	0.379	0.337	0.311	0.278	0.245	0.210	0.163	0.142
$\alpha_B^{(2loop)}$	0.64	0.359	0.328	0.293	0.27	0.244	0.217	0.188	0.15	0.131
$\alpha_s^{(2loop)}$		0.73	0.404	0.33	0.293	0.256	0.229	0.19	0.149	0.131

Table 2: $\alpha_B(Q^2)$ computed numerically to all orders within minimal model, Eq.(8), with $\Lambda = 0.37$ GeV, $\Delta = 1.5$, $m = M_0 = 1$ GeV and in extended model, Eqs. (3), (21), with $\eta = 3$, $\tilde{C} = 1.253$ reproducing all known coefficients $\beta_n, n = 0, \dots, 3$ in $\overline{MS}$ scheme for $n_f = 0$. The last two lines refer to the two-loop approximation of α_B and α_s (in SPT) with the same parameters.

References

- [1] S.P. Booth et al., Phys. Lett. **B294** (1992) 385;
G.S. Bali and K. Schilling, Phys. Rev. **D47** (1993) 661;
G.S. Bali, Phys. Lett. **B460** (1999) 170.
- [2] Yu.A. Simonov, Yad. Fiz. **58** (1995) 113; JETP Lett. **57** (1993) 513, hep-ph/9311247;
Yu.A. Simonov, in: Lecture Notes in Physics, v. 479, p. 139 (1996).

- [3] A.M. Badalian, Yu.A. Simonov, Phys. Atom. Nucl. **60** (1997) 630.
- [4] A.M. Badalian, V.L. Morgunov, Phys. Rev. **D60** (1999) 116008.
- [5] A.M. Badalian and B.L.G. Bakker, Phys. Rev. **D62** (2000) 094031.
- [6] A.M. Badalian and D.S. Kuzmenko, hep-ph/0104097.
- [7] G. 't Hooft, Nucl. Phys. **B72** (1974) 461;
E. Witten, Nucl. Phys. **B156** (1979) 269.
- [8] L.F. Abbot, Nucl. Phys. **185** (1981) 189.
- [9] Review of Particle Physics, Europ. Phys. Journ. **C15** (2000) 229.
- [10] Yu.A. Simomov, hep-ph/0109081; Phys. Atom. Nucl. (in press).
- [11] E.C. Poggio, H.R. Quinn and S. Weinberg, Phys. Rev. **D13** (1976) 1958.
- [12] A. Radyushkin, JINR preprint E-2-82-159 (1982) and hep-ph/9901228;
B. Schrempp and F. Schrempp, Z.Phys. **C6** (1980) 7;
N.V. Krasnikov and A.A. Pivovarov, Phys. Lett. **116 B** (1982) 168;
see also D.V. Shirkov hep-ph/0012283.
- [13] F.J. Yndurain, The Theory of Quark and Gluon Interactions, 3d edition, Springer- Verlag, Berlin-Heidelberg, 1999.
- [14] A.C. Mattingly and P.M. Stevenson, Phys. Rev. **D49** (1994) 437;
Yu.L. Dokshitzer, hep-ph/0106348; hep-ph/9812252.